

OBSERVATIONAL EVIDENCE FOR MASSIVE BLACK HOLES IN THE CENTERS OF ACTIVE GALAXIES

J. M. Moran, L. J. Greenhill

Harvard-Smithsonian Center for Astrophysics

and J. R. Herrnstein

National Radio Astronomy Observatory

ABSTRACT

Naturally occurring water vapor maser emission at 1.35 cm wavelength provides an accurate probe for the study of accretion disks around highly compact objects, thought to be black holes, in the centers of active galaxies. Because of the exceptionally fine angular resolution, 200 microarcseconds, obtainable with very long baseline interferometry, accompanied by high spectral resolution, $< 0.1 \text{ km s}^{-1}$, the dynamics and structures of these disks can be probed with exceptional clarity. The data on the galaxy NGC 4258 are discussed here in detail. The mass of the black hole binding the accretion disk is $3.9 \times 10^7 M_{\odot}$. Although the accretion disk has a rotational period of about 800 years, the physical motions of the masers have been directly measured with VLBI over a period of a few years. These measurements also allow the distance from the earth to the black hole to be estimated to an accuracy of 4 percent. The status of the search for other maser/black hole candidates is also discussed.

1. INTRODUCTION

The observational evidence for the existence of supermassive black holes (10^6 – 10^9 times the mass of the sun, $M_{\odot}$) in the centers of active galaxies has been accumulating at an ever accelerating pace for the last few decades (e.g., Rees 1998; Blandford & Gehrels 1999). Seyfert (1943) first drew attention to a group of galaxies with unusual excitation conditions in their nuclei indicative of energetic activity. Among the twelve galaxies in his list was NGC 4258, which is the subject of much of this paper. Such galaxies, now known as galaxies with active galactic nuclei (AGN), have grown in membership and importance. Ironically, NGC 4258 no longer belongs to the class of Seyfert galaxies by modern classification standards (Heckman 1980), but it is still considered to have a mildly active galactic nucleus. Meanwhile, the study of AGN has become a major field in modern astrophysics. In the 1960s, galaxies with AGN were discovered with intense radio emission arising from jets of relativistic particles often extending far beyond the optical boundaries of the host galaxy. The central engine, the source of energy that powers such jets and other phenomena in the centers of galaxies, has long been ascribed to black holes (e.g., Salpeter 1964; Blandford & Rees 1992). There are two sources of energy for these phenomena:

the gravitational energy from material falling onto the black hole and the spin energy of the black hole itself (Blandford & Znajek 1977).

The direct evidence for black holes in AGN has come principally from observations of the motions of gas and stars in the extended environments of black holes. In the optical and infrared domains, the evidence for black holes from stellar measurements comes from an analysis of the velocity dispersion of stars as a function of distance from the dynamical centers of galaxies. In the case of our own Galactic center, the proper motions (angular velocities in the plane of the sky) of individual stars can be measured. These data show that there is a mass of about $2.6 \times 10^6 M_\odot$ within a volume of radius 0.01 pc (Genzel et al. 1997; Ghez et al. 1998). In addition, measurements by the Hubble Space Telescope of the velocity field of hydrogen gas in active galaxies indicate the presence of massive centrally condensed objects. Reviews of these data have been written by Faber (1999), Ho (1999), Kormendy and Richstone (1995), and others.

In the X-ray portion of the spectrum, there is compelling evidence for black holes in AGN from the detection of the highly broadened iron $K\alpha$ line at 6.4 keV. The line is broadened by the gravitational redshift of gas as close as 3 Schwarzschild radii from the black hole. An example of an iron line profile in the galaxy MCG-6-30-15 is shown in Figure 1 (Tanaka et al. 1995). The linear extent of the emission region cannot be determined directly by the X-ray telescope, so it is not possible to estimate directly the mass of the putative black hole. Detailed analysis of the line profile suggests that the black hole is spinning (e.g., Bromley, Miller, & Pariev 1998).

In the radio regime, a new line of inquiry has given unexpectedly clear and compelling evidence for black holes: the discovery of water masers orbiting highly massive and compact central objects. With the aid of very long baseline interferometry (VLBI), which provides angular resolution as fine as 200 microarcseconds (μas) at a wavelength of 1.3 cm and spectral resolution of 0.1 km s^{-1} or less, the structure of accreting material around these central objects can be studied in detail. This paper describes the observations and the significance of these measurements of water masers in AGN. We begin with a brief description of cosmic masers and the interferometric techniques used to observe them.

2. COSMIC MASERS

Intense maser action in cosmic molecular clouds was discovered in 1965 (Weaver et al. 1965) from observations of OH, and later from H₂O, SiO, and CH₃OH. In the case of water vapor, the commonly observed masers emit in the $6_{16}-5_{23}$ transition at 22235 MHz (1.35 cm wavelength). Most masers have been found to be associated with one of two types of objects: newly formed stars or evolved stars (e.g., Elitzur 1992; Reid & Moran 1988). Although very distinct, they share the characteristic of having envelopes of outflowing gas and dust (silicate material). The pump source in all cases is thought to come in the form of either shock waves or infrared radiation. More recently, masers have been found in the spiral arms of nearby galaxies and in AGN.




Fig. 1.— The X-ray spectrum of the galaxy MCG-6-30-15, observed by the Japanese ASCA satellite. The top panel shows the total spectrum with a model of the continuum emission fitted to the data outside the range of 5–7 keV. The bottom panel shows the residuals, which reveal a broad spectral feature attributed to the Fe K α line at 6.4 keV. The line has a width of 100,000 km s⁻¹. The most extreme redshifted part is thought to arise from gas at a radius of about 3 Schwarzschild radii. (From Tanaka et al. 1995)

Cosmic masers are similar to their laboratory counterparts on earth in that their intense radiation is produced by population inversion. However, cosmic masers are one-pass amplifiers and have little temporal or spatial coherence. The intensity of cosmic masers varies, often erratically, on timescales of hours to years. The underlying electric field is a Gaussian random process (Moran 1981). In an unsaturated maser (in astronomical terminology), the pumping is sufficiently strong that the microwave intensity does not affect the level populations, and the intensity increases exponentially through the masing medium. The input signal can be either a background source or the maser's own spontaneous emission. In a saturated maser, one pump photon is needed for each microwave photon, and in a one-dimensional maser medium where beaming can be neglected, the intensity increases linearly with distance. Maser emission is expected to be beamed. Most masers are thought to be saturated, and this condition requires the least pump power. However, this assertion is difficult to prove observationally.

Consider a simple geometry for a maser, a filamentary tube, shown in Figure 2. The boundaries of the filament can be defined in terms of either gas density or the region where the line-of-sight velocity is constant to within the thermal line width. The gas in the tube is predominantly molecular hydrogen, with trace amounts of water vapor (about one part in 10⁵)




Fig. 2.— A cartoon of a simple filamentary maser. The pump energy can be supplied by either shock waves or radiation. Pump cycles usually involve excitation to infrared rotational levels, followed by de-excitation to the upper level of the maser transition. In order to avoid thermalization, the infrared photons emitted during the pump cycle must escape from the masing medium, which favors a geometry that is thin in at least one dimension. In a filamentary geometry, the masing medium is defined either by the physical extent of the masing gas or by the volume over which the gas is coherent, that is, where the variation in the velocity projected along the line of sight is less than the thermal line width. The maser emission is beamed into a cone of angular opening equal to d/L . If the maser amplifies a background source, the radiation will be beamed in the forward direction. If it amplifies its own spontaneous emission, then beamed maser emission will emerge from both ends, and weaker emission from the sidewalls.

and other constituents. If this maser medium is saturated, then the luminosity is given by

$$L = h\nu n \Delta P V , \quad (1)$$

where h is Planck's constant, ν is the frequency, n is the population density in pumping level, ΔP is the differential pump rate per molecule, and V is the volume of the masing cloud. This is the most luminosity a maser of given pump rate and volume can produce. The emission will be beamed along the major axis of the filament into an angle

$$\beta \cong \frac{d}{L} , \quad (2)$$

where d is the cross-sectional diameter, and L is the length of the filament. The beam angle and the length of the maser are not directly observable. Since the volume of the maser is approximately $d^2 L$, the observed flux density from a maser beamed toward the earth is

$$F_\nu = \frac{1}{2} h\nu n \frac{\Delta P}{\Delta \nu} \frac{V}{D^2 \Omega} = \frac{1}{2} h\nu n \frac{\Delta P}{\Delta \nu} \frac{L^3}{D^2} , \quad (3)$$

where $\Delta \nu$ is the line width, D is the distance between the maser and the observer, and Ω is the beam angle of the emission, $\sim \beta^2$. The maximum allowable hydrogen number density is about 10^{10} molecules per cubic centimeter, above which the maser levels become thermalized by collisions. This maximum allowable density ($\rho_c = 3 \times 10^{-13} \text{ g cm}^{-3}$, an important parameter in much of the




Fig. 3.— The top panel shows the maser spectrum as first discovered in NGC 4258 at velocities near the systemic velocity of the galaxy (Claussen, Heiligman, & Lo 1984). The lower spectrum shows the observations of Nakai, Inoue, & Miyoshi (1993) over a much broader velocity range. The velocities are computed from the Doppler effect and are based on a rest frequency of 22235.080 MHz. The effects of the motions of the earth and sun with respect to the local standard of rest have been removed. The bars indicate velocity ranges of emission.

following discussion) along with the maximum allowable pump rate, which equals the Einstein A-coefficient for infrared transitions linking the maser levels, limit the luminosity of a maser of given volume.

Water masers outside our Galaxy were first discovered in the spiral arms of the nearby galaxy M33 by Churchwell et al. (1977). Their properties were found to be similar to masers found in Galactic star-forming regions. Much more luminous water masers were found in the AGN associated with NGC 4945 and the Circinus galaxy by dos Santos & Lepine (1979) and Gardner & Whiteoak (1982), respectively. Claussen, Heiligman, & Lo (1984) and Claussen & Lo (1986) conducted surveys and found five additional masers associated with AGN, including the one in NGC 4258 (see Figure 3). They suggested that these masers might arise in gas associated with dust-laden molecular tori that had been proposed to surround black holes by Antonucci & Miller (1985). Nakai, Inoue, & Miyoshi (1993), with a powerful new spectrometer of 16,000 channels spanning a velocity range of 3000 km s^{-1} , observed NGC 4258 and discovered satellite line clusters offset from the systemic velocity by about $\pm 1000 \text{ km s}^{-1}$, which are shown in Figure 3 (see also Miyoshi 1999).

The compelling reason that the radio emission from the water vapor transition arises from the maser process is straightforward. A typical example of a maser line from a small part of




Fig. 4.— An expanded view of the spectral feature near 1306 km s^{-1} in NGC 4258, which is typical of emission from an isolated maser component (see Figure 3). The line width is characteristic of gas at 300 K, but the intensity corresponds to that of a blackbody at an equivalent temperature greater than 10^{14} K .

the spectrum of NGC 4258 is shown in Figure 4. The line width is about 1 km s^{-1} , the thermal broadening expected for a gas cloud at 300 K. However, the angular size determined by radio interferometry is less than $100 \mu\text{as}$, implying that the equivalent blackbody temperature must exceed 10^{14} K . The exceedingly high brightness of the radiation is the principal evidence for the maser process. (Typical molecular lines from molecular clouds have velocities of several tens of km s^{-1} — due to thermal and turbulent broadening — and brightness temperatures of less than 100 K.) Cosmic masers produce very bright spots of radiation but have little else in common with terrestrial masers. It is difficult to use masers to determine physical conditions (e.g., temperature, density) in molecular clouds because of the complexity of the maser process. However, as compact sources of narrowband radiation, masers are ideal probes of the dynamics of their environment.

3. VLBI

The most important tool for the study of the angular structure of masers is very long baseline interferometry (VLBI). Signals from a maser, or from other bright compact radio sources, are converted to a low-frequency baseband and recorded in digital format on magnetic tape at Nyquist sampling rates of up to about 10^8 samples per second at widely separated telescopes that operate independently. They form a radio version of the classical Michelson stellar interferometer, whose coherence is maintained by the use of atomic frequency standards to preserve the signal phase and timing (see Figure 5). The received signals (which are proportional to the incident electric fields)




Fig. 5.— A block diagram of a two-element very long baseline interferometer. It operates as a coherent interferometer. An atomic frequency standard (F.S.) controls the phase of the local oscillator signal at each telescope used to convert the radio frequency signal to a video band for recording on magnetic tape in digital form (without square-law detection) and sampled at the appropriate Nyquist rate. On playback, one of the signals is delayed by τ to compensate for the differential propagation time from the source to the antennas. The signals are correlated and Fourier transformed to produce cross-power spectra, or correlated power as a function of frequency and time.

from an array of two or more telescopes are cross-correlated pairwise to form cross-correlation functions. Taking advantage of the earth's rotation, the spatial cross-correlation function of the incident electromagnetic field, or visibility, can be measured over a wide range of projected baseline vectors. The image and fringe visibility functions are related through a Fourier transform (see Thompson, Moran, & Swenson 1986). The temporal Fourier transform of the cross-correlation function gives the cross-power spectrum of the radiation, or visibility as a function of frequency, so that images at different frequencies can be obtained. The intrinsic angular resolution, θ , of a multielement interferometer is $0.7\lambda/B$, where λ is the wavelength, and B is the longest baseline length. For water vapor, $\lambda = 1.35$ cm, and B is typically 6000 km, which gives a resolution of $200 \mu\text{as}$. The spectral resolution available is typically about 15 KHz, or about a fifth of the line widths (see Figure 4). In maser sources, one spectral feature at a particular frequency or velocity can be used as a phase reference for the interferometer, and all other phases referred to it. With this technique, the coherence time of the interferometer can be extended indefinitely, and the relative positions of masers with respect to the reference feature can be measured to a small fraction of the fringe spacing, or intrinsic resolution. The relative position of an unresolved maser




Fig. 6.— The distribution of the ten elements of the Very Long Baseline Array (VLBA). This network is often augmented with other radio telescopes such as the Very Large Array (a 27-element array operating as a phased array), shown here with the symbol Y, and the 100 m telescope of the Max Planck Institute for Radio Astronomy near Bonn, Germany.

component can be measured to an accuracy of about

$$\Delta\theta = \frac{1}{2} \frac{\theta}{SNR}, \quad (4)$$

where SNR is the signal-to-noise ratio.

4. THE STUDY OF NGC 4258

The imaging of the maser in NGC 4258 was one of the first projects undertaken by a dedicated VLBI system known as the Very Long Baseline Array (VLBA) (see Figure 6) in the spring of 1994. Previous VLBI measurements of the systemic features had shown that they arose from an elongated structure with a velocity gradient along the major axis, highly suggestive of a rotating disk seen edge-on (Greenhill et al. 1995a). The VLBA measurements of all the maser components provide convincing evidence for a rotating disk around a massive central object.

The basic observational results on NGC 4258 obtained over the past few years can be summarized as follows (see Table 1 for a list of parameters):

1. The masers appear to trace a highly elongated, although slightly curved, structure (Figure 7). The high-velocity, redshifted and blueshifted features are offset in position on the left and right sides of the systemic features, respectively. The velocities of the high-velocity features as a function of impact parameter (position along the major axis of the distribution) follow the prediction of Kepler's third law of orbital motion. The systemic features show linear dependence with impact parameter (Miyoshi et al. 1995).

TABLE 1

PARAMETERS OF MOLECULAR DISK TRACED BY WATER VAPOR MASERS IN NGC 4258^a

Inner radius, R_i	0.14 pc (3.9 mas)
Outer radius, R_o	0.28 pc (8.0 mas)
Inner rotation velocity, $v_\phi(R_i)$	1100 km s ⁻¹
Outer rotation velocity, $v_\phi(R_o)$	770 km s ⁻¹
Inner rotation period	800 yrs
Outer rotation period	2200 yrs
Position angle of disk (at 3.9 mas radius)	80°
Inclination angle	98°
Position velocity slope	282 km s ⁻¹ mas ⁻¹
Central mass, M	$3.9 \times 10^7 M_\odot$
Disk mass, M_d	$< 10^6 M_\odot$
Central mass density, uniform distribution	$> 4 \times 10^9 M_\odot \text{ pc}^{-3}$
Central mass, Plummer distribution	$> 10^{12} M_\odot \text{ pc}^{-3}$
Centripetal acceleration, systemic features	9.3 km s ⁻¹ yr ⁻¹
Centripetal acceleration, high-velocity features	$\leq 0.8 \text{ km s}^{-1} \text{ yr}^{-1}$
Disk systemic velocity ^b , v_0	476 km s ⁻¹
Galactic systemic velocity (optical) ^{b,c}	472 km s ⁻¹
Radial drift velocity, v_R	$< 10 \text{ km s}^{-1}$
Thickness of disk, H	$< 0.0003 \text{ pc}$
Maser beam angle, β	8°
Disk–galaxy angle ^d	119°
Apparent maser luminosity	150 L _⊙
Model luminosity ^e	11 L _⊙
Distance, D	$7.2 \pm 0.3 \text{ Mpc}$

^aBased on the distance estimate of 7.2 Mpc.^bRadio definition, with respect to the local standard of rest. To convert to heliocentric velocity (radio), subtract 8.2 km s⁻¹; to convert to heliocentric (optical), subtract 7.5 km s⁻¹.^cFrom Cecil, Wilson, & Tully (1992).^dAngle between the spin axis of the molecular disk and the spin axis of the galaxy.^eRadiation into a zone within $\pm 4^\circ$ of the plane of the disk.

2. The distribution of maser features in the direction normal to the major axis is too small to be measured at present (see Figure 7). The upper limit on the ratio of thickness to radius of the disk is 0.0025 (Moran et al. 1995).




Fig. 7.— Top: Image of the maser emission from the nucleus of NGC 4258. The ticks on the axes are in milliarcseconds. One milliarcsecond corresponds to 0.035 pc, or 1.1×10^{17} cm, at a distance of 7.2 Mpc. Bottom: The line-of-sight velocities of the masers versus position along the major axis. The curved portions of the plot precisely follow a Keplerian dependence. Data from January 1995 (top) and April 1994 (bottom).

3. The upper limit of any toroidal component of the magnetic field in the masers, derived from searches for Zeeman splitting in the line at 1306 km s^{-1} , is less than 300 mG (Herrnstein et al. 1998a).
4. The accelerations (i.e., the linear drift in the line-of-sight velocity with time) of the systemic features are about $9 \text{ km s}^{-1} \text{ yr}^{-1}$ (Haschick, Baan, & Peng 1994; Greenhill et al. 1995b; Nakai et al. 1995). The high-velocity features that have been tracked have accelerations in the range $\pm 0.8 \text{ km s}^{-1} \text{ yr}^{-1}$ (Bragg et al. 1999).
5. The high-velocity features show no proper motions with respect to a fixed-velocity component in the systemic range (Herrnstein 1996). The systemic features show proper motions of about $32 \mu\text{as yr}^{-1}$ (Herrnstein et al. 1999).
6. There is an elongated continuum radio source, which appears to be a jet emanating from the black hole position, parallel to the axis of rotation (Herrnstein et al. 1997). There is no 1.35 cm wavelength emission from the position of the black hole (Herrnstein et al. 1998b).

There is virtually no doubt that the masers trace a very thin disk in nearly perfect Keplerian motion. Five of six phase-space parameters have been measured for each maser spot, two spatial coordinates and three velocity coordinates. The missing coordinate is the position along the line of sight, which must be inferred from the constraint provided by Kepler's third law.

The approximate placement of the masers in the disk can be understood by considering a simple thin, flat disk viewed edge-on. In this case the line-of-sight velocity, v_z , of a maser will be given by

$$v_z - v_0 = \sqrt{\frac{GM}{R}} \sin \phi, \quad (5)$$

where v_0 is the line-of-sight velocity of the central object (i.e., the systemic velocity), G is the gravitational constant, R is the distance of a maser component from the black hole, and ϕ is the azimuth angle in the disk, measured from the line between the black hole and the observer. If the disk were randomly filled with observable masers, one might expect to see a velocity position diagram as shown in Figure 8. The linear boundaries of the distribution are populated by masers at the inner and outer edges of the annular disk. The masers on the curved boundaries lie on the midline, where $\phi = 90^\circ$. Hence, the masers in NGC 4258 have a very specific distribution: the high-velocity masers lie close to the midline, and the systemic masers lie within a narrow range of radii. From Equation 5, the radius of a particular maser can be determined as

$$R = (GM)^{\frac{1}{3}} \left[\frac{b}{v_z - v_0} \right]^{\frac{2}{3}}, \quad (6)$$

where b is the projected distance on the sky along the major axis from the center of the disk ($\sin \phi = b/R$). Similarly, positional offsets from the midline, z , of the high-velocity features can be determined by deviations from a Keplerian curve; that is,

$$z = \sqrt{R^2 - b^2}. \quad (7)$$

There is a two-fold ambiguity in the z component of the position for the edge-on disk case. Unambiguous estimates of the positions of the high-velocity features have been derived from the accelerations by Bragg et al. (1999), who showed that the masers lie within 15 degrees of the midline.

An expanded plot of the Keplerian part of the velocity curve is shown in Figure 9. The data fit a Keplerian curve to an accuracy of about 3 km s^{-1} , or less than 1 percent of the rotation speed. However, there are noticeable deviations from a perfect fit. The estimate of the central mass of the disk derived from this data depends on the distance to the maser, and has a value of $3.9 \times 10^7 M_\odot$ for a distance of 7.2 Mpc. This mass corresponds to an Eddington luminosity (where radiation pressure from Thomson scattering would balance gravity) of $5 \times 10^{45} \text{ erg s}^{-1}$. Since the total electromagnetic emission appears to be less than $10^{42} \text{ erg s}^{-1}$, the system is highly sub-Eddington.




Fig. 8.— Top: A cartoon model of a flat annular disk viewed edge-on, with randomly distributed maser sources. Bottom: Each maser will appear as a point within the “bow tie” boundary in the plot of line-of-sight velocity versus impact parameter. The curved portion of the boundary is populated by masers located along the midline, the diameter perpendicular to the line of sight. This is a pure Keplerian curve, because the velocity vectors lie along the line of sight. Masers at a fixed radius will appear along a straight line. The steep and shallow lines correspond to masers on the inner and outer annular boundaries of the disk.

Since this binding mass must lie inside the inner radius of the maser disk, the mass density, assuming a spherical mass distribution, is at least $4 \times 10^9 \text{ M}_\odot \text{ pc}^{-3}$ ($3 \times 10^{-13} \text{ g cm}^{-3}$). It is unlikely that this mass is in the form of a dense star cluster (Maoz 1995). The average density of stars in the solar neighborhood is about $1 \text{ M}_\odot \text{ pc}^{-3}$, and the density of the densest known star cluster is about $10^5 \text{ M}_\odot \text{ pc}^{-3}$. A star cluster will have a mass distribution that decreases monotonically with radius. In order not to disrupt the Keplerian curve, the core mass for a reasonable distribution must have a peak density of at least $1 \times 10^{12} \text{ M}_\odot \text{ pc}^{-3}$. A cluster of massive stars at this density would evaporate from gravitational interactions on a timescale short with respect to the age of the galaxy, while a cluster of low-mass stars would destroy itself from collisions over a similar timescale. Hence, it is unlikely that the central mass is in the form of a star cluster (see also Begelman & Rees 1978). The best explanation is that the central object is a supermassive black hole, with a Schwarzschild radius (R_S) of about $1.2 \times 10^{13} \text{ cm}$. Hence, the masers are distributed




Fig. 9.— The magnitude of the line-of-sight velocities of the masers, relative to the systemic velocity, versus distance from the dynamical center of NGC 4258. The filled circles are the redshifted masers, and the squares are the blueshifted masers. (Data from April 1994)

in a zone between 40,000 and 80,000 R_S . Because the maser clouds are so far from the event horizon, deviations of their motions from the predictions of Newtonian mechanics are small. The gravitational redshift and transverse Doppler shift are about 4 km s^{-1} (detectable), the expected Lense-Thirring precession (see below) is less than about 3° over the maser annulus (possibly detectable), and the apparent shift of the maser positions due to gravitational bending is about $0.1 \mu\text{as}$ (undetectable).

The disk is remarkably thin. In a disk supported against gravity by pressure (hydrostatic equilibrium), the density distribution is expected to have a Gaussian profile with a thickness, H , given by the relation

$$H/R = (c_s^2 + v_a^2)^{1/2} / v_\phi, \quad (8)$$

where c_s is the sound speed and v_a is the Alfvén speed, which characterize thermal and magnetic support pressure, respectively, and v_ϕ is the Keplerian rotational speed. Since $H/R < 0.0025$, the quadrature sum of the sound speed and Alfvén speed is less than 2.5 km s^{-1} . The upper limit on the magnetic field of 300 mG suggests that the Alfvén speed, $B/\sqrt{4\pi\rho}$, where ρ is the density, is less than 3 km s^{-1} for $\rho = \rho_c$ (the critical density for quenching maser emission). If the support were completely due to thermal pressure, the temperature would be less than 1000 K.

A proper determination of the positions of the masers on the disk requires that the warp and the inclination of the disk to the line of sight be taken into account. An example of such a disk, slightly warped (in position angle only) and slightly inclined to the line of sight, that fits the maser distribution in position and velocity is shown in Figure 10.




Fig. 10.— The warped annular disk (wire mesh) modeled to the maser positions, velocities, and accelerations (adapted from Herrnstein, Greenhill, & Moran 1996). The black dot in the center marks the dynamical center of the disk. The continuum emission at 1.3 cm is shown in the shaded gray contours. The southern jet may be weaker than the northern jet because of thermal absorption in the disk. The lack of emission at the position of the black hole places constraints on any coronal or advection zone surrounding the black hole (Herrnstein et al. 1998b).

The distance to the maser of 7.2 ± 0.3 Mpc was determined from analysis of the proper motions and accelerations of the systemic features (Herrnstein et al. 1999). Fifteen features were tracked over a period of two years to an accuracy of $0.5\text{--}10 \mu\text{as}$ in relative position and 0.4 km s^{-1} in velocity. The distance estimate is based on simple geometric considerations. The Keplerian curve of the high-velocity masers gives the mass function $GM \sin^2 i/D$, where i is the inclination of the disk to the line of sight. The radius, R , of the systemic masers (in angular units) is determined from Equation (6), based on the slope of the velocity versus impact parameter curve shown in Figure 7. This fixes the angular velocity, v_ϕ , of the systemic masers under the assumption that the orbits are circular. The expected accelerations and proper motions of the systemic features are v_ϕ^2/R and v_ϕ/D , respectively. The assumption that the orbits are circular is reasonable on theoretical grounds because of viscous relaxation and on observational grounds because the continuum emission arises close to the center of symmetry of the maser distribution.

The distance to the 15 Cepheid variables in NGC 4258 has been estimated to be 8.1 ± 0.8 Mpc (Maoz *et al.* 1999). The statistical component of the error is 0.4 Mpc and the systematic error associated with the calibration of the Cepheid distance scale is 0.7 Mpc. The discrepancy between the two distance measurements to NGC 4258 may have cosmological implications (Paczynski 1999).

5. INTERESTING UNANSWERED QUESTIONS

1. What is the rate of radial inflow of material through the disk?

The accretion rate of material onto the black hole is an important parameter that affects our understanding of radiation processes around the black hole. The maser data provide some

information about the accretion rate. Key issues are the long timescale needed for material to flow from the disk to the black hole and the assumption that the masers trace all the disk material. It is useful to first estimate the mass of the disk. If we account for systematic effects in addition to the random scatter of 3 km s^{-1} , the deviation from Keplerian motion due to the finite mass of the disk, Δv_ϕ , is less than about 10 km s^{-1} over the radius of the disk. This limits the mass of the disk to less than about $2M\Delta v_\phi/v_\phi$, or about $10^6 M_\odot$. The density of the molecular gas must be less than ρ_c (10^{10} hydrogen molecules per cubic centimeter). Since $H/R < 0.0025$, the upper limit on mass is $10^5 M_\odot$. If, in addition, the disk is stable against the effects of self-gravity (Toomre 1964; Binney & Tremaine 1987), then the mass of the disk must be less than $M(H/R)$, or about $10^5 M_\odot$.

The mass accretion rate of a disk in steady state is given by

$$\dot{M} = 2\pi R \Sigma v_R, \quad (9)$$

where Σ is the surface density of the disk, and v_R is radial drift velocity, which depends on the viscosity of the disk. Unfortunately, v_R is only weakly constrained by the observations (i.e., the possible difference between the optical and radio systemic velocities) to be $< 10 \text{ km s}^{-1}$. This provides a crude limit on the accretion rate of $100 M_\odot \text{ yr}^{-1}$. To further constrain the mass accretion rate requires an estimate of the viscosity of the disk. In the standard model of a thin, viscous accretion disk, as formulated by Shakura and Sunyaev (1973), v_R can be written (see Frank, King, & Raine 1992) as

$$v_R = \alpha v_\phi \left(\frac{H}{R} \right)^2, \quad (10)$$

where α is the dimensionless viscosity parameter ($0 \leq \alpha \leq 1$). The observational limit on the ratio H/R implies that $v_R < 0.006\alpha \text{ km s}^{-1}$. With the limit on mass given by the deviation from Keplerian motion, the accretion rate is less than $10^{-1}\alpha M_\odot \text{ yr}^{-1}$. The infall time from the masing region is R/v_R , which from Equation (10) can be written as

$$T \sim \frac{1}{\alpha} \left(\frac{c}{c_S} \right)^2 \left(\frac{R_S}{c} \right) \left(\frac{R}{R_S} \right)^{\frac{1}{2}}. \quad (11)$$

For NGC 4258, with $\alpha = 0.1$ and $c_S = 2.5 \text{ km s}^{-1}$, $T = 10^{16} \text{ s}$, or about $3 \times 10^8 \text{ yrs}$.

From the magnetic field limit and the assumption of equipartition of magnetic and thermal energy, the upper limit on $\dot{M}$ is also $10^{-1}\alpha M_\odot \text{ yr}^{-1}$. If the maser density is the maximum allowable value, ρ_c , and the maser traces all the material in the disk, then the limit on disk thickness leads to an upper limit on $\dot{M}$ of $10^{-2}\alpha M_\odot \text{ yr}^{-1}$. Detailed theoretical modeling can give estimates for the accretion rate. For example, a model in which the cause of the outer radial cutoff in maser emission is attributed to the transition from molecular to atomic gas leads to an estimate of $10^{-4}\alpha M_\odot \text{ yr}^{-1}$ (Neufeld & Maloney 1995). Gammie, Narayan, & Blandford (1999) favor an accretion rate of $10^{-1}\alpha M_\odot \text{ yr}^{-1}$, based on an analysis of the continuum radiation spectrum.

If the accretion rate is high, then the relative weakness of the continuum radiation may be due to the process of advection (Gammie, Narayan, & Blandford 1999). On the other hand, if the accretion rate is low, then the weak emission is due to the dearth of infalling material. In this case the gravitational power in the accretion flow may be insufficient to power the jets.

2. What is the form and origin of the warp?

The form of the warp is difficult to determine precisely, because the filling factor of the masers in the disk is so small. Better measurements of the positions and directions of motion of the high-velocity features are key to defining the warp more accurately.

The cause of the warp is unknown, but several suggestions have been put forward. Papaloizou, Terquem, & Lin (1998) show that the warp could be produced by a binary companion orbiting outside the maser disk. Its mass would need to be comparable to the mass of the disk ($< 10^6 M_\odot$). Alternatively, radiation pressure from the central source will produce torques on a slightly warped disk and will cause the warp to grow (Maloney, Begelman, & Pringle 1996). Finally, it is conceivable that in the absence of other torques, the observed warp is due to the Lense-Thirring effect. A maximally rotating black hole will cause a precession of a nonaligned orbit (weak field limit) of

$$\Omega_{LT} = \frac{2G^2 M^2}{c^3 R^3}, \quad (12)$$

which can be rewritten in terms of the Schwarzschild radius as

$$\Omega_{LT} = \frac{1}{2} \frac{c}{R_S} \left(\frac{R_S}{R} \right)^3. \quad (13)$$

At the inner radius of the disk ($R/R_S = 40,000$), the precession amounts to $3 \times 10^{-17} \text{ s}^{-1}$. This precession is very small but might be significant over the lifetime of the disk. Equation (11) suggests that the lifetime might be 10^{16} s , which would produce a differential precession of about 10° across the radius of the disk. If the axis of the disk is inclined to the axis of the black hole, then the viscosity of the disk is expected to twist the plane of the innermost part of the disk to the equatorial plane of the black hole (Bardeen & Petterson 1975; Kumar & Pringle, 1985).

3. Do the water masers trace the whole disk?

The inner and outer radii of the observed masers are undoubtedly due to excitation conditions in the maser. In the vertical direction it is also possible that the masers form in a thin region within a thicker disk with atomic and ionized components. It has also been proposed that the high-velocity features are not indicative of a warped disk but trace material that has been blown off a flat disk (Kartje, Konigl, & Elitzur 1999). These proposals are difficult to test.

4. What are the physical properties of the maser spots?

The spectrum of the maser has many discrete peaks which correspond to spots of maser




Fig. 11.— A cartoon of the molecular accretion disk in NGC 4258. The period of rotation at the inner edge of the disk is about 800 years. A particular systemic maser is visible for about 10 years as its radiation beam, estimated to have a width of about 8° , sweeps over the earth. The high-velocity features may be visible for a substantially longer time. (Adapted from Greenhill et al. 1995a)

emission on the sky. The success of measuring proper motions and accelerations of these masers suggests that they correspond to discrete condensations or density-enhanced regions in the disk. A cartoon of the blobs in a disk is shown in Figure 11. The blobs in front of the black hole may be visible because they amplify emission from the central region. No masers have been seen on the backside of the disk. On the other hand, the high-velocity masers have no continuum emission to amplify, and we may only see the ones near the midline, where the gradient in the line-of-sight velocity is small. Blobs in the rest of the disk may be radiating in directions away from the earth. The clumpiness of the medium allows us to track the individual masers. If the appearance of spots is due to blobs, or density enhancements in the disk, then the minimum gradient condition would not seem to be necessary. However, intense high-velocity maser spots may occur when blobs at the same velocity line up to form two-stage masers (Deguchi & Watson 1989). This situation forms a highly beamed maser, like the filamentary maser described in Section 2. The probability of realizing this situation is greatest along the midline, where the velocity gradient is smallest. All evidence suggests that the masers arise from discrete physical condensations. There have been several suggestions that the apparent motions of the maser spots may be due to a phase effect (e.g., a spiral density wave moving through the disk, Maoz & McKee 1998), but there is no

observational evidence for this.

6. MASERS IN OTHER AGN

At this time (early 1999), 22 masers have been detected among about 700 galaxies searched (e.g., Braatz, Wilson, & Henkel 1997). A list of these galaxies with masers is given in Table 2.

TABLE 2
KNOWN AGN WITH WATER MASERS

Galaxy	Distance (Mpc)	Flux Density (Jy)	Milliarcsecond Structure	Disk
M 51	3	0.2	...	...
NGC 4945	3.7	4	yes	yes
Circinus	4	4	yes	yes
NGC 4258	7.2	4	yes	yes
NGC 1386	12	0.9	yes	maybe
NGC 3079	16	6	yes	maybe
NGC 1068	16	0.6	yes	yes
NGC 1052	20	0.3	yes	no
NGC 613	20	0.1?	...	...
NGC 5506	24	0.6	...	...
NGC 5347	32	0.1	...	...
NGC 3735	36	0.2	...	...
IC 2560	38	0.4	yes	...
NGC 2639	44	0.1	...	...
NGC 5793	50	0.4	...	...
ESO 103-G035	53	0.7	...	...
Mrk 1210	54	0.2	...	...
IRAS F01063-8034	57	0.2	...	...
Mrk 1	65	0.1	...	...
NGC 315	66	0.05	...	...
IC 1481	83	0.4	...	...
IRAS F22265-1826	100	0.3	yes	no

NOTE— Extragalactic masers outside AGN are found in NGC 253, M 82, M 33, IC 342, LMC, and SMC.

The yield rate of detections is only about 3 percent. The major reason for this paucity is probably that the maser disks can only be seen if they are edge-on to the line of sight. If the typical beam angle, β , is 8° , as in NGC 4258, then the probability of seeing a maser is about equal to $\sin \beta$, or 8 percent. Braatz, Wilson, & Henkel (1997) have shown that most of the known masers are associated with Seyfert II galaxies or LINERs where the accretion disks are thought to be edge-on to the earth.

It is difficult to make VLBI measurements on masers weaker than about 0.5 Jy because of the need to detect the maser within the coherence time of the interferometer. Nine masers have been studied with VLBI. Four of these show strong evidence of disk structure, and two more show probable disk structure. The properties of these masers are listed in Table 3. Unfortunately, none

TABLE 3
WATER MASERS WITH RESOLVED STRUCTURE

Masers Without Obvious Disk Structure							
Galaxy	D Mpc	v_0 km s^{-1}	Δv km s^{-1}	ΔR pc	Comment	Reference	
IRAS 22265 (S0)	100	7570	150	2.4	messy	Greenhill et al. 1999a	
NGC 1052 (E4)	20	1490	100	0.6	“jet”	Claussen et al. 1998	
IC 2560	38	2900	30	0.2	velocity gradient	Nakai et al. 1998	
Masers With Disk Structure							
Galaxy	D Mpc	v_ϕ km s^{-1}	R_i/R_o pc	M $10^6 \text{ M}_\odot$	ρ $10^7 \text{ M}_\odot \text{ pc}^{-3}$	L_x $10^{42} \text{ erg s}^{-1}$	Reference
NGC 4258	7	1100	0.13/0.26	35	400	0.04	Miyoshi et al. 1995
NGC 1068	15	330	0.6/1.2	17	3	40	Greenhill & Gwinn 1997
Circinus	4	230	0.08/0.8	1	40	40	Greenhill et al. 1999b
NGC 4945	4	150	0.2/0.4	1	2	1	Greenhill et al. 1997
NGC 1386	12	100	–/0.7	2	4	0.02	Braatz et al. 1999
NGC 3079	16	150	–/1.0	1	0.2	0.02	Trotter et al. 1998, Satoh et al. 1998

D = distance, v_0 = systemic velocity, Δv = velocity range, ΔR = linear extent, v_ϕ = rotational velocity, R_i/R_o = inner/outer radius of disk, M = central mass, ρ = central mass density, L_x = X-ray luminosity.

of these masers show the simple, well-defined structure that would make them useful for precise study of the physical properties of accretion disks around black holes.

7. SUMMARY

The measurements of the positions and velocities of the masers in the nucleus of NGC 4258 offer compelling evidence for the existence of a supermassive black hole and provide the first direct image of an accretion disk within $10^5 R_S$ of the black hole. Much more can be learned from this system. A measurement of the disk thickness is important and may require higher signal-to-noise ratios than are achievable currently or VLBI measurements from space. Measurement of the continuum spectrum from the central region is very important to the understanding of the radiation process. Detection of radio emission would require instruments of higher sensitivity. Continued measurements over time of the positions and velocities of the masers will refine the estimates of their proper motions and accelerations, and this will better define the shape of the disk. It is even conceivable that the radial drift velocity will be detected. This work will benefit immensely from new instruments that are in the planning stage for centimeter wavelength radio astronomy. These include the enhanced Very Large Array, the Square Kilometer Array, and space VLBI missions such as ARISE.

We thank Adam Trotter and Ann Bragg for helpful discussions.

REFERENCES

- Antonucci, R. R. J. & Miller, J. S. 1985, *Astrophys. J.*, **297**, 621–632.
- Bardeen, J. M. & Petterson, J. A. 1975, *Astrophys. J.*, **195**, L65–L67.
- Begelman, M. C. & Rees, M. J. 1978, *Mon. Not. R. Astr. Soc.*, **185**, 847–859.
- Binney, J. & Tremaine, S. 1987, *Galactic Dynamics* (Princeton: Princeton Univ. Press).
- Blandford, R. D. & Gehrels, N. 1999, *Physics Today*, **52**, 40–46.
- Blandford, R. D. & Rees, M. 1992, in *Testing the AGN Paradigm*, ed. S. S. Holt, S. G. Neff, & C. M. Urry (New York: American Inst. of Physics), 3–19.
- Blandford, R. D. & Znajek, R. L. 1977, *Mon. Not. R. Astr. Soc.*, **179**, 433–456.
- Bragg, A. E., Greenhill, L. J., Moran, J. M., & Henkel, C. 1999, *Astrophys. J.*, submitted.
- Braatz, J. A., Wilson, A. S., & Henkel, C. 1997, *Astrophys. J. Supp.*, **110**, 321–346.
- Braatz, J. A., et. al. 1999, *Astrophys. J.*, in preparation.
- Bromley, B. C., Miller, W. A., & Pariev, V. I. 1998, *Nature*, **391**, 54–56.

- Cecil, G., Wilson, A. S., & Tully, R. B. 1992, *Astrophys. J.*, **390**, 365–377.
- Churchwell, E., Witzel, A., Huchtmeier, W., Pauliny-Toth, I., Roland, J., & Sieber, W. 1977, *Astr. Astrophys.*, **54**, 969–971.
- Claussen, M. J., Diamond, P. J., Braatz, J. A., Wilson, A. S., & Henkel, C. 1998, *Astrophys. J.*, **500**, L129–L132.
- Claussen, M. J., Heiligman, G. M., & Lo, K. Y. 1984, *Nature*, **310**, 298–300.
- Claussen, M. J. & Lo, K.-Y. 1986, *Astrophys. J.*, **308**, 592–599.
- Deguchi, S. & Watson, W. D. 1989, *Astrophys. J.*, **340**, L17–L20.
- dos Santos, P. M. & Lepine, J. R. D. 1979, *Nature*, **278**, 34–35.
- Elitzur, M. 1992, *Astronomical Masers* (Dordrecht: Kluwer).
- Faber, S. M. 1999, in Proceedings of the 32nd COSPAR Meeting, *The AGN-Galaxy Connection*, ed. H R. Schmitt, L. C. Ho, & A. L. Kinney (Advances in Space Research), in press.
- Frank, J., King, A., & Raine, D. 1992, *Accretion Power in Astrophysics* (Cambridge: Cambridge Univ. Press).
- Gammie, C. F., Narayan, R., & Blandford, R. 1999, *Astrophys. J.*, **516**, 177–186.
- Gardner, F. F. & Whiteoak, J. B. 1982, *Mon. Not. R. Astr. Soc.*, **201**, 13p–15p.
- Genzel, R., Eckart, A., Ott, T., & Eisenhauer, F. 1997, *Mon. Not. R. Astr. Soc.*, **291**, 219–234.
- Ghez, A. M., Klein, B. L., Morris, M., & Becklin, E. E. 1998, *Astrophys. J.*, **509**, 678–686.
- Greenhill, L. J., et al. 1999a, in preparation.
- Greenhill, L. J., et al. 1999b, in preparation.
- Greenhill, L. J. & Gwinn, C. R. 1997, *Astrophys. Space Sci.*, **248**, 261–267.
- Greenhill, L. J., Henkel, C., Becker, R., Wilson, T. L., & Wouterloot, J. G. A. 1995b, *Astr. Astrophys.*, **304**, 21–33.
- Greenhill, L. J., Jiang, D. R., Moran, J. M., Reid, M. J., Lo, K. Y., & Claussen, M. J. 1995a, *Astrophys. J.*, **440**, 619–627.
- Greenhill, L. J., Moran, J. M., & Herrnstein, J. R. 1997, *Astrophys. J.*, **481**, L23–L26.
- Haschick, A. D., Baan, W. A., & Peng, E. W. 1994, *Astrophys. J.*, **437**, L35–L38.
- Heckman, T. M. 1980, *Astr. Astrophys.*, **87**, 152–164.
- Herrnstein, J. R. 1996, Ph.D. thesis, Harvard University.

- Herrnstein, J. R., Greenhill, L. J., & Moran, J. M. 1996, *Astrophys. J.*, **468**, L17–L20.
- Herrnstein, J. R., Greenhill, L. J., Moran, J. M., Diamond, P. J., Inoue, M., Nakai, N., & Miyoshi, M. 1998b, *Astrophys. J.*, **497**, L69–L73.
- Herrnstein, J. R., Moran, J. M., Greenhill, L. J., Blackman, E. G., & Diamond, P. J. 1998a, *Astrophys. J.*, **508**, 243–247.
- Herrnstein, J. R., Moran, J. M., Greenhill, L. J., Diamond, P. J., Inoue, M., Nakai, N., Miyoshi, M., Henkel, C., & Riess, A. 1999, *Nature*, **400**, 539–541.
- Herrnstein, J. R., Moran, J. M., Greenhill, L. J., Diamond, P. J., Miyoshi, M., Nakai, N., & Inoue, M. 1997, *Astrophys. J.*, **475**, L17–L20.
- Ho, L. C. 1999, in *Observational Evidence for Black Holes in the Universe*, ed. S. K. Chakrabarti (Dordrecht: Kluwer), 157–186.
- Kartje, J. F., Konigl, A., & Elitzur, M. 1999, *Astrophys. J.*, **513**, 180–196.
- Kumar, S. & Pringle, J. E. 1985, *Mon. Not. R. Astr. Soc.*, **213**, 435–442.
- Kormendy, J. & Richstone, D. 1995, *Ann. Rev. Astr. Astrophys.*, **33**, 581–624.
- Maloney, P. R., Begelman, M. C., & Pringle, J. E. 1996, *Astrophys. J.*, **472**, 582–587.
- Maoz, E. 1995, *Astrophys. J.*, **455**, L131–L134.
- Maoz, E., *et al.* 1999, *Nature*, **401**, 351–354.
- Maoz, E. & McKee, C. 1998, *Astrophys. J.*, **494**, 218–235.
- Miyoshi, M. 1999, in *Observational Evidence for Black Holes in the Universe*, ed. S. K. Chakrabarti (Dordrecht: Kluwer), 141–156.
- Miyoshi, M., Moran, J. M., Herrnstein, J. R., Greenhill, L. J., Nakai, N., Diamond, P. D., & Inoue, M. 1995, *Nature*, **373**, 127–129.
- Moran, J. 1981, *Bull. Am. Astron. Soc.*, **13**, 508.
- Moran, J. M., Greenhill, L. J., Herrnstein, J. R., Diamond, P. D., Miyoshi, M., Nakai, N., & Inoue, M. 1995, *Proc. Nat. Acad. Sci., USA*, **92**, 11427–11433.
- Nakai, N., Inoue, M., Hagiwara, Y., Miyoshi, M., & Diamond, P. J. 1998, in *Radio Emission from Galactic and Extragalactic Compact Sources*, ed. J. A. Zensus, G. B. Taylor, & J. M. Wrobel (San Francisco: ASP), 237–238.
- Nakai, N., Inoue, M., Miyazawa, K., Miyoshi, M., & Hall, P. 1995, *Pub. Astron. Soc. Jap.*, **47**, 771–799.
- Nakai, N., Inoue, M., & Miyoshi, M. 1993, *Nature*, **361**, 45–47.

- Neufeld, D. & Maloney, P. R. 1995, *Astrophys. J.*, **447**, L17–L20.
- Paczynski, B. 1999, *Nature*, **401**, 331–332.
- Papaloizou, J. C. B., Terquem, C., & Lin, D. N. C. 1998, *Astrophys. J.*, **497**, 212–226.
- Reid, M. J. & Moran, J. M. 1988, in *Galactic and Extragalactic Radio Astronomy*, 2d ed., ed. G. L. Verschuur & K. I. Kellermann (New York: Springer-Verlag), 255–294.
- Rees, M. 1998, in *Black Holes and Relativistic Stars*, ed. R. M. Wald (Chicago: Univ. Chicago Press), 79–101.
- Salpeter, E. E. 1964, *Astrophys. J.*, **140**, 796–800.
- Satoh, S., Inoue, M., Nakai, N., Shibata, K. M., Kamenoi, S., Migenes, V., & Diamond, P. J. 1998, in *Highlights of Astronomy*, **11b**, ed. J. Andersen (Dordrecht: Kluwer), 972–973.
- Seyfert, C. 1943, *Astrophys. J.*, **97**, 28–40.
- Shakura, N. I. & Sunyaev, R. A. 1973, *Astr. Astrophys.*, **24**, 337–355.
- Tanaka, Y., et al. 1995, *Nature*, **375**, 659–661.
- Thompson, A. R., Moran, J. M., & Swenson, G. W. 1986, *Interferometry and Synthesis in Radio Astronomy* (New York: Wiley Interscience).
- Toomre, A. 1964, *Astrophys. J.*, **139**, 1217–1238.
- Trotter, A. S., Greenhill, L. J., Moran, J. M., Reid, M. J., Irwin, J. A., & Lo, K. Y. 1998, *Astrophys. J.*, **495**, 740–748.
- Weaver, H., Williams, D. R. W., Dieter, N. H., & Lum, W. T. 1965, *Nature*, **208**, 29–31.

QUESTIONS AND ANSWERS

Q: What other measurements can one do with masers to try to get sensitivity to the general relativistic effects or to get information from distances closer to the mass concentration?

A: We have searched for masers in NGC 4258 that are closer to the central mass than 40,000 Schwarzschild radii and have found none. The reason for the absence of masers closer to the center is unclear. Masers are very sensitive to local conditions such as temperature, density, and pump power, which may depend on the geometry of the warp. Water masers arise in neutral media at temperatures below a few thousand degrees, so it is unrealistic to expect to find masers very close to the Schwarzschild radius.

One possible general relativity effect might be the deflection of radiation from masers on the far side of the accretion disk (i.e., a distortion of the image of the back side of the accretion disk). However, no such masers have been found.